

Fast Facts

	<i>Symptoms</i>	<i>Incubation Period</i>	<i>Location</i>
Malaria	Fever, chills, headache, muscle aches, tiredness, nausea, vomiting, diarrhea. Can cause bleeding, kidney failure, seizures, mental confusion, jaundice, coma, death in severe cases.	10 - 30 Days Can be up to 1 year later	Worldwide, but largest number of cases in sub-Saharan Africa
Dengue Fever	High fever, severe headache, severe pain behind the eyes, joint pain, muscle/bone pain, rash, mild bleeding (nose/gums, easily bruised).	4 - 7 Days	Central and northern part of South America, parts of Africa and South Asia
West Nile	No symptoms (70-80% of people infected), fever, headache, body aches, joint pain, vomiting, diarrhea, rash, fatigue/weakness.	2 – 6 Days Can be 2 – 14 Days	United States, Europe, Middle East, Africa, West Asia
Chikungunya	Fever, joint pain, headache, muscle pain, joint swelling, rash, fatigue/weakness.	3 – 7 Days	United States (including Alaska), central and northern part of South America, Asia, Italy, France, parts of Africa
Zika	Fever, rash, joint pain, muscle pain, headache, conjunctivitis (red eyes).	Unknown	Central and northern part of South America, Oceania/Pacific Islands, Cape Verde (Africa)

©2016 Demystifying Your Health. All rights reserved.

Demystifying Your Health does not provide medical advice, diagnosis or treatment.